



Turbulent Nature of the Quasicontinuous Exhaust Regime for Fusion Plasmas

Paper Rehearsal


Kaiyu Zhang*, Wladimir Zholobenko, Andreas Stegmeir, Michael Faitsch, Konrad Eder, Christoph Pitzal, Frank Jenko, and ASDEX Upgrade Team





The Quasicontinuous Exhaust regime (QCE)

- ELM-free H-mode with large λ_q ,
high n_{sep} in favor of detachment
- Experimental Observation:
 1. Separatrix spanning **Quasi-coherent mode (QCM)**
 2. Intermittent filaments (**plasma blobs**) in the SOL
- First-principles understanding
→ Turbulence simulations using GRILLIX



[M. Faitsch, Nuclear Materials and Energy, 2021]



GRILLIX: two-fluid turbulence codes for edge & SOL

Full-f, global

$n_0 + n_1, T_{e0} + T_{e1}, T_{i0} + T_{i1}$ evolve together $\rightarrow n_1/n_0 \gtrsim 1$

\rightarrow coefficients (resistivity, viscosity, ionization rate ...)

Locally-field-aligned method

[A. Stegmeir, Comput. Phys. Comm. 2023]

Parallel: field aligned

Perpendicular: cartesian grid

\rightarrow High resolution across the separatrix & X-point

$$\Delta l_{\perp} \sim \rho_s$$



Electromagnetics

Turbulence-driven magnetic fluctuations

$$\mathbf{B}_1 \neq 0$$

[K. Zhang, Nucl. Fusion 2024]

[K. Zhang, Comput. Phys. Comm. 2024]





Phenomenology & Validation

Dynamics

- **Quasi-coherent mode (QCM)**

$$k_{\text{pol}} \rho_s = 0.033$$


[J. Kalis, Nucl. Fusion, 2024]


- **Blobs (filaments)**

$$v_{\text{blob}}^r \sim 1 \text{ km/s}$$

[M. Griener, Nucl. Materials and Energy, 2020]

$$L_{\perp} \sim 1 \text{ cm} \quad L_{\parallel} > 10 \text{ m}$$





Phenomenology & Validation


Dynamics

- **Quasi-coherent mode (QCM)**
 $k_{\text{pol}}\rho_s = 0.033$
 [J. Kalis, Nucl. Fusion, 2024]
 - **Blobs (filaments)**
 $v_{\text{blob}}^r \sim 1 \text{ km/s}$
 [M. Griener, Nucl. Materials and Energy, 2020]
- $L_{\perp} \sim 1 \text{ cm}$ $L_{\parallel} > 10 \text{ m}$

Plasma profiles

Global, full-f, **first-principles**


- 3-moment neutral gas model with recycling boundary
- Landau fluid closure for q_{\parallel}
 Neoclassical correction for $\eta_{\text{vis},\parallel}$
- **No free parameters**





Fluctuation statistics across the separatrix


1. QCM: strong fluctuations within $\pm 2.5\sigma$ (continuous) at sep.
2. Voids $< -2.5\sigma$ (intermittent) inside sep.
3. Blobs $> 2.5\sigma$ (intermittent) outside sep.



Two separatrix-spanning modes underlying QCE



[J. R. Myra, Phys. Plasmas, 2005]



Kinetic ballooning mode (KBM)

'Winner' at outboard-mid plane

$$\omega_r \sim \omega_{kbm} \sim \frac{\omega_{*i}}{2}$$

$$\alpha_{\phi,n} \approx \pi$$

Resistive X-point mode (RXM)

'Loser' relegated near X-point

$$\omega_r \sim \omega_{resi}$$

$$\alpha_{\phi,n} \approx 0$$


Quasi-coherent mode in experiment

[J. Kalis, Nucl. Fusion 2024]

[G. Grenfell, Nucl. Fusion 2024]

1. Separatrix: QCM is dominated by KBM
2. Near SOL: QCM undergoes $\alpha_{\phi,n} \rightarrow \pi/2$, where KBM & RXM's eigenfunctions merge, which ejects blobs.

Interchange for blobs [D. A. D'Ippolito, Phys. Plasmas, 2011]




Blobs



Mechanism for combining high-confinement & broad SOL fall-off




AUG [M. Faitsch, Nucl. Materials and Energy, 2021]

1. Observe **decoupled decay lengths** where $\alpha_{\phi,n} = \pi/2$ and QCM ejects blobs
2. Enhanced SOL transport is carried by intermittent fluctuations (**blobs**)
3. Pedestal-foot transport is dominated by **KBM**



Transport properties at the pedestal foot

Experimentally, QCM avoids ELM, but can be very close to Ideal ballooning threshold





Mechanism for mesoscopic radial extent of QCM: electromagnetic self-organization

QCM resides inside a **broader** E_r well.

AUG [G. F. Harrer, Phys. Rev. Lett. 2022]


1. **High- n_{sep}** is a key lever for controlling QCM.

AUG [M. Faitsch, Nucl. Fusion, 2023]
C-Mod [M.A. Miller, Nucl. Fusion, 2025]

2. **Maxwell stress** depletes zonal flow. Without it, E_r & $\partial_r p_i$ evolve deeper, QCM vanishes.

$$\frac{\partial}{\partial t} \frac{\partial \langle nu^\vartheta \rangle}{\partial \varrho} = \underbrace{-\frac{\partial^2}{\partial \varrho^2} \langle nu^e u^\vartheta \rangle}_{\text{Reynolds stress}} + \underbrace{\frac{\hat{\Omega}_i}{e} \frac{\mu_0}{B_{ax}^2} \frac{\partial}{\partial \varrho^2} \langle B_1^e B_1^\vartheta \rangle}_{\text{Maxwell stress}} + \text{others}$$

$$E_r = \frac{\partial_r p_i}{en} + u_\varphi B_\theta - u_\theta B_\varphi$$





Mechanism for preventing violent transport


- **Finite Larmor radius effects**, sustaining $\alpha_{\phi,n} \sim \pi$

Linear KBM

$$\frac{d}{d\theta} (1 + (s\theta - \alpha_m \sin \theta)^2) \frac{d}{d\theta} \frac{\tilde{A}_{\parallel}}{K_{\parallel}} = \left[C_0 + C_2 \omega^2 + C_1 \omega + \frac{C_{-1}}{\omega - \omega_0} \right] \frac{\tilde{A}_{\parallel}}{K_{\parallel}}$$


ballooning angle magnetic shear $\propto \frac{\partial \beta}{\partial r}$ **Ideal Ballooning Mode** **Finite Larmor radius** **Magnetic Drift Resonance**
 $\alpha_{\phi,n} = \pi/2$ $\alpha_{\phi,n} = \pi/2 \rightarrow \pi$ reduce unstable threshold

Turbulence



Reduce FLR

Consequences
 Heat flux $\times 10$
 Pedestal drops





Conclusions

- **Validation** against a wide set of experiment metrics:
mean profiles, fluctuation spectra, mode structure
- **Two separatrix-spanning modes**
Kinetic ballooning mode, **KBM** (outboard midplane)
Resistive X-point mode, **RXM** (X-point)
- **Quasi-coherent mode**: a self-organized mesoscopic **KBM**,
with Finite Larmor radius effects and Maxwell stress regulating its transport
- **Blobs**: ejected by interchange dynamics during the interaction between **RXM** and **KBM**
- The blob-dominated SOL temperature fall-off is well decoupled from
the pedestal-foot gradient set by the QCM.